

# Engineering Casimir interactions with metamaterials

Diego A. R. Dalvit  
Theoretical Division  
Los Alamos National Laboratory



# Collaborators

## LANL

Peter Milonni

Felipe da Rosa

Francesco Intravaia

Antoniette Taylor

John O'Hara

## Sandia

Paul Davids

Igal Brener

Maarten de Boer

Stephen Howell

## MIT

Steven Johnson

Alejandro Rodriguez

Alex McCauley

# Outline of this Talk


## ■ Casimir forces and metamaterials

- What is a metamaterial?
- Proposals for Casimir force manipulation with metamaterials
- Is Casimir repulsion possible?

## ■ Some very new results (paper under review)

- Stay till the end....

# Metamaterials and Casimir






# Effects of materials

The Lifshitz formula: Lifshitz (1956)

$$\frac{F}{A} = 2\hbar \operatorname{Im} \int_0^\infty \frac{d\omega}{2\pi} \int \frac{d^2 \mathbf{k}_\parallel}{(2\pi)^2} K_3 \operatorname{Tr} \frac{\mathbf{R}_1 \cdot \mathbf{R}_2 e^{2iK_3 d}}{1 - \mathbf{R}_1 \cdot \mathbf{R}_2 e^{2iK_3 d}}$$

$$K_3 = \sqrt{\omega^2/c^2 - k_\parallel^2}$$



Reflection matrices (Fresnel formulas for isotropic media):

$$r^{\text{TM},\text{TM}}(\omega, \mathbf{k}_\parallel) = \frac{\epsilon(\omega)K_3 - \sqrt{\epsilon(\omega)\mu(\omega)\omega^2/c^2 - k_\parallel^2}}{\epsilon(\omega)K_3 + \sqrt{\epsilon(\omega)\mu(\omega)\omega^2/c^2 - k_\parallel^2}}$$

$$r^{\text{TE},\text{TE}} = r^{\text{TM},\text{TM}} \quad \text{with} \quad \epsilon \leftrightarrow \mu$$


Relevant frequencies:

$$\left. \begin{array}{ll} \epsilon(\omega) \xrightarrow{\omega \gg \Omega_p} 1 & \Rightarrow r^{p,p} \approx 0 \text{ (Transparent plates)} \\ \omega \gg c/d & \Rightarrow e^{2iK_3 d} \approx 0 \text{ (Fast oscillations)} \end{array} \right\} \Rightarrow F \approx 0$$

$$0 \leq \omega \leq \min\{\Omega_p, c/d\}$$

# Going to imaginary frequencies

$$\frac{F}{A} = 2\hbar \int_0^\infty \frac{d\xi}{2\pi} \int \frac{d^2\mathbf{k}_\parallel}{(2\pi)^2} K_3 \text{Tr} \frac{\mathbf{R}_1 \cdot \mathbf{R}_2 e^{-2K_3 d}}{1 - \mathbf{R}_1 \cdot \mathbf{R}_2 e^{-2K_3 d}}$$



Kramers-Kronig (causality) relations:

$$\epsilon(i\xi) = 1 + \frac{2}{\pi} \int_0^\infty \frac{\omega \epsilon''(\omega)}{\omega^2 + \xi^2} d\omega$$

$$\mu(i\xi) = 1 + \frac{2}{\pi} \int_0^\infty \frac{\omega \mu''(\omega)}{\omega^2 + \xi^2} d\omega$$

Dominant frequencies below the near-infrared/optical region of the EM spectrum (gaps  $d = 200\text{-}1000$  nm)

The important message is that Casimir is a broad-band frequency phenomenon

# The sign of the Casimir force

$$\frac{F}{A} = 2\hbar \int_0^\infty \frac{d\xi}{2\pi} \int \frac{d^2\mathbf{k}_\parallel}{(2\pi)^2} K_3 \text{Tr} \frac{\mathbf{R}_1 \cdot \mathbf{R}_2 e^{-2K_3 d}}{1 - \mathbf{R}_1 \cdot \mathbf{R}_2 e^{-2K_3 d}}$$

The sign of the force is directly connected to the **sign of the product of the reflection coefficients** on the two plates, **evaluated at imaginary frequencies**. As a rule of thumb, we have (p=TE,TM)

$$R_1^p(i\xi) \cdot R_2^p(i\xi) > 0 \quad (\forall \xi \leq c/d) \Rightarrow \text{Attraction}$$

$$R_1^p(i\xi) \cdot R_2^p(i\xi) < 0 \quad (\forall \xi \leq c/d) \Rightarrow \text{Repulsion}$$

In terms of permittivities and permeabilities:


$$\begin{array}{l} \epsilon_a(i\xi) \gg \epsilon_b(i\xi) \\ \mu_b(i\xi) \gg \mu_a(i\xi) \end{array} \longrightarrow \text{Repulsion}$$

# Ideal attraction-repulsion

## ■ Ideal attractive limit

Casimir (1948)

$$\frac{F}{A} = +\frac{\pi^2}{240} \frac{\hbar c}{d^4}$$




# Ideal attraction-repulsion

## ■ Ideal attractive limit

Casimir (1948)


$$\frac{F}{A} = + \frac{\pi^2}{240} \frac{\hbar c}{d^4}$$



## ■ Ideal repulsive limit

Boyer (1974)

$$\frac{F}{A} = - \frac{7}{8} \frac{\pi^2}{240} \frac{\hbar c}{d^4}$$




# Ideal attraction-repulsion

## ■ Ideal attractive limit

Casimir (1948)


$$\frac{F}{A} = + \frac{\pi^2}{240} \frac{\hbar c}{d^4}$$



## ■ Ideal repulsive limit

Boyer (1974)

$$\frac{F}{A} = - \frac{7}{8} \frac{\pi^2}{240} \frac{\hbar c}{d^4}$$



## ■ Real repulsive limit


Casimir repulsion is associated with strong electric-magnetic interactions. However, natural occurring materials do NOT have strong magnetic response in the optical region, i.e.  $\mu = 1$

# Ideal attraction-repulsion

## ■ Ideal attractive limit

Casimir (1948)


$$\frac{F}{A} = + \frac{\pi^2}{240} \frac{\hbar c}{d^4}$$



## ■ Ideal repulsive limit

Boyer (1974)

$$\frac{F}{A} = - \frac{7}{8} \frac{\pi^2}{240} \frac{\hbar c}{d^4}$$



## ■ Real repulsive limit

Casimir repulsion is associated with strong electric-magnetic interactions. However, natural occurring materials do NOT have strong magnetic response in the optical region, i.e.  $\mu = 1$

→ **Metamaterials**

# Quantum levitation with MMs?

Physicists have 'solved' mystery of levitation - Telegraph

<http://www.telegraph.co.uk/news/main.jhtml?xml=/news/2007/08/0...>

RSS | Britain's No.1 quality newspaper website | Make us your homepage

Telegraph.co.uk

Home News Sport Business Travel Jobs Motoring Property SEARCH Go Car site Web

News home  
Law reports  
Obituaries  
Picture galleries

Announcements  
Arts  
Blogs  
Comment  
Crossword  
Dating  
Digital Life  
Earth  
Expert  
Family  
Fantasy Games  
Fashion  
Features  
Food & Drink  
Football  
Gardening  
Health  
Horoscopes  
My Telegraph  
Obituaries  
Promotions  
Science  
Sudoku  
Sunday Telegraph  
Telegraph e-paper  
Telegraph magazine  
Telegraph offers  
Telegraph PM  
Weather  
Your Money  
Your view

NEWS SERVICES  
BlackBerry service  
Desktop alerts  
Email services  
Home delivery  
Mobile  
Photographs  
Podcasts  
RSS feeds  
Weekly Telegraph

FEATURE FOCUS

How green is your home?

## Physicists have 'solved' mystery of levitation

By Roger Highfield, Science Editor  
Last Updated: 1:41pm BST 08/08/2007

Levitation has been elevated from being pure science fiction to science fact, according to a study reported today by physicists.

In earlier work the same team of theoretical physicists showed that invisibility cloaks are feasible.

Now, in another report that sounds like it comes out of the pages of a Harry Potter book, the University of St Andrews team has created an 'incredible levitation effects' by engineering the force of nature which normally causes objects to stick together.

Professor Ulf Leonhardt and Dr Thomas Philbin, from the University of St Andrews in Scotland, have worked out a way of reversing this phenomenon, known as the Casimir force, so that it repels instead of attracts.

Their discovery could ultimately lead to frictionless micro-machines with moving parts that levitate. But they say that, in principle at least, the same effect could be used to levitate bigger objects too, even a person.

advertisement

The Casimir force is a



In theory the discovery could be used to levitate a person

consequence of quantum mechanics, the theory that describes the world of atoms and subatomic particles that is not only the most successful theory of physics but also the most baffling.

The force is due to neither electrical charge or gravity, for example, but the fluctuations in all-pervasive energy fields in the intervening empty space between the objects and is one reason atoms stick together, also explaining a "dry glue" effect that enables a gecko to walk across a ceiling.

Now, using a special lens of a kind that has already been built, Prof Ulf Leonhardt and Dr Thomas Philbin report in the New Journal of

Physics they can engineer the Casimir force to repel, rather than attract.

Because the Casimir force causes problems for nanotechnologists, who are trying to build electrical circuits and tiny mechanical devices on silicon chips, among other things, the team believes the feat could initially be used to stop tiny objects from sticking to each other.

Prof Leonhardt explained, "The Casimir force is the ultimate cause of friction in the nano-world, in particular in some microelectromechanical systems.

Such systems already play an important role - for example tiny mechanical devices which triggers a car airbag to inflate or those which power tiny 'lab on chip' devices used for drugs testing or chemical analysis.

Micro or nano machines could run smoother and with less or no friction at all if one can manipulate the force." Though it is possible to levitate objects as big as humans, scientists are a long way off developing the technology for such feats, said Dr Philbin.

The practicalities of designing the lens to do this are daunting but not impossible and levitation "could happen over quite a distance".

Prof Leonhardt leads one of four teams - three of them in Britain - to have put forward a theory in a peer-reviewed journal to achieve invisibility by making light waves flow around an object - just as a river flows undisturbed around a smooth rock.

Print Email this story

Post this story to: del.icio.us | Digg | Newsvine | NowPublic | Reddit

Most Viewed

REGION BY REGION

Harry Potter star 'disgusted' at friend's sex act  
Will President Bush bomb Iran?  
Threat to take new-born over emotional abuse  
Russian bombers launch missiles over Arctic  
M1 coach crash driver 'drink driving' arrest






“In theory the discovery could be used to levitate a person”





# Metamaterials

- Artificial structured composites with designer electromagnetic properties
- MMs are strongly anisotropic, dispersive, magneto-dielectric media.

- Negative refraction** Veselago (1968), Smith et al (2000)
- Perfect lens** Pendry (2000)
- Cloaking** Smith et al (2007)




THz MMs: eg split ring resonators




$$\epsilon, \mu < 0$$

Optical MMs: eg nano-pillars



200nm




# Effective medium approximation

We want to compute the Casimir force between a metallic plate and a MM.  
Let us assume a metallic plate is reasonably well described by a Drude response

$$\epsilon_1(\omega) = 1 - \frac{\Omega_e^2}{\omega^2 + i\gamma_e\omega} \quad \mu_1 = 1$$

For the MM the optical response is not so simple.....



In the effective medium approximation (EMA) one describes the MM with an effective electric permittivity and an effective magnetic permeability. This is an approximation valid when the MM is probed at wavelengths much larger than the average distance between the constituent “particles” of the MM.



# EMA: Electric response






Close to resonance, the optical response can be modeled by a Drude-Lorentz permittivity

$$\varepsilon(\omega) = 1 - \frac{\omega_p^2 - \omega_0^2}{\omega^2 - \omega_0^2 + i\omega\Gamma}$$



# EMA: Magnetic response

$$\mu_{eff} = 1 - \frac{\frac{\pi r^2}{a^2}}{1 + \frac{2\sigma i}{\omega r \mu_0} - \frac{3}{\pi^2 \mu_0 \omega^2 C r^3}}$$



J.B. Pendry *et al.*, *IEEE Trans. Microwave Tech.* **47**, 2075 (1999).

# EMA: Drude-Lorentz responses

Close to the resonance, both  $\epsilon(\omega)$  and  $\mu(\omega)$  can be modeled by Drude-Lorentz formulas

$$\epsilon_{\alpha}(\omega) = 1 - \frac{\Omega_{E,\alpha}^2}{\omega^2 - \omega_{E,\alpha}^2 + i\Gamma_{E,\alpha}\omega}$$

$$\mu_{\alpha}(\omega) = 1 - \frac{\Omega_{M,\alpha}^2}{\omega^2 - \omega_{M,\alpha}^2 + i\Gamma_{M,\alpha}\omega}$$

Typical separations

$$d = 200 - 1000 \text{ nm}$$




Infrared-optical frequencies

$$\Omega/2\pi = 5 \times 10^{14} \text{ Hz}$$


Drude metal (Au)

$$\Omega_E = 9.0 \text{ eV} \quad \Gamma_E = 35 \text{ meV}$$



Metamaterial

$$\text{Re } \epsilon_2(\omega) < 0 \quad \text{Re } \mu_2(\omega) < 0$$



$$\Omega_{E,2}/\Omega = 0.1 \quad \Omega_{M,2}/\Omega = 0.3$$


$$\omega_{E,2}/\Omega = \omega_{M,2}/\Omega = 0.1$$

$$\Gamma_{E,2}/\Omega = \Gamma_{M,2}/\Omega = 0.01$$

# Attraction-repulsion crossover

Drude metal (Au)


Drude metal (Au)



Only attraction

Metamaterial


Metamaterial



Only attraction

Drude metal (Au)

Metamaterial




Repulsion-attraction



# Drude background


In some metallic-based MMs, there is a net conductivity due to the metallic structure, like the fishnet design on the right.



$$\epsilon(\omega) = 1 - f \frac{\Omega_D^2}{\omega^2 - i\omega\gamma_D} - (1 - f) \frac{\Omega_e^2}{\omega^2 - \omega_e^2 + i\gamma_e\omega}$$

$f$  : filling factor

$$\mu(\omega) = 1 - \frac{\Omega_m^2}{\omega^2 - \omega_m^2 + i\gamma_m\omega}$$







A Drude background is detrimental for Casimir force reduction or repulsion, since it results in an electric response much stronger than the magnetic one

$$\epsilon_2(i\xi) \gg \mu_2(i\xi)$$

# Complementary SRRs + Drude

SRRs structures provide an opportunity to avoid the large Drude background, since they can be built in two natural complementary ways





# Optical anisotropy


In an anisotropic medium, the constitutive relations between  $\mathbf{E}$ ,  $\mathbf{D}$ ,  $\mathbf{B}$ , and  $\mathbf{H}$  are more involved:

$$\mathbf{D} = \epsilon \cdot \mathbf{E} \quad \mathbf{H} = \mu^{-1} \cdot \mathbf{B}$$

due to the tensorial nature of the permittivity and permeability

$$\epsilon = \begin{bmatrix} \epsilon_{11} & \epsilon_{12} & \epsilon_{13} \\ \epsilon_{21} & \epsilon_{22} & \epsilon_{23} \\ \epsilon_{31} & \epsilon_{32} & \epsilon_{33} \end{bmatrix} \quad \mu = \begin{bmatrix} \mu_{11} & \mu_{12} & \mu_{13} \\ \mu_{21} & \mu_{22} & \mu_{23} \\ \mu_{31} & \mu_{32} & \mu_{33} \end{bmatrix}$$


due to the tensorial nature of the permittivity and permeability



Examples of uniaxial  
anisotropy in stacked MMs



# Anisotropy: Uniaxial MMs



$$\epsilon_{ij} = \begin{bmatrix} \epsilon_{xx} & 0 & 0 \\ 0 & \epsilon_{xx} & 0 \\ 0 & 0 & \epsilon_{zz} \end{bmatrix} ; \quad \mu_{ij} = \begin{bmatrix} \mu_{xx} & 0 & 0 \\ 0 & \mu_{xx} & 0 \\ 0 & 0 & \mu_{zz} \end{bmatrix}$$

Anisotropy produces polarization mixing  
(non-diagonal reflection matrices)


$$\epsilon_{xx}(\omega) = \epsilon_{yy}(\omega) = 1 - (1 - f_x) \frac{\Omega_{e,x}^2}{\omega^2 - \omega_{e,x}^2 + i\gamma_{e,x}\omega} - f_x \frac{\Omega_{D,x}^2}{\omega^2 + i\gamma_{D,x}\omega},$$

$$\epsilon_{zz}(\omega) = 1 - (1 - f_z) \frac{\Omega_{e,z}^2}{\omega^2 - \omega_{e,z}^2 + i\gamma_{e,z}\omega} - f_z \frac{\Omega_{D,z}^2}{\omega^2 + i\gamma_{D,z}\omega},$$

$$\mu_{xx}(\omega) = \mu_{yy}(\omega) = 1 - \frac{\Omega_{m,x}^2}{\omega^2 - \omega_{m,x}^2 + i\gamma_{m,x}\omega},$$

$$\mu_{zz}(\omega) = 1 - \frac{\Omega_{m,z}^2}{\omega^2 - \omega_{m,z}^2 + i\gamma_{m,z}\omega}$$

Anisotropy does not help repulsion



(Rosa, DD, Milonni, PRA 2008)

# EMA: correct model for $\mu$

Drude-Lorentz for permeability is wrong. The correct expression that results in EMA from Maxwell's equations is


$$\mu_{\text{eff}}(\omega) = 1 - f \frac{\omega^2}{\omega^2 - \omega_m^2 + 2i\gamma_m\omega} \quad (\text{Pendry 1999})$$

The appearance of the  $\omega^2$  factor in the numerator is very important:

Although close to the resonance this behaves in the same way as the Drude-Lorentz EMA permeability, it has a completely different low-frequency behavior




$$\mu_{\text{eff}}(i\xi) < 1 < \epsilon_{\text{eff}}(i\xi)$$

**No Casimir repulsion!**



# Other Casimir MMs: chirality

The chirality of a MM is defined by the chirality of its unit cell



In a chiral medium, the constitutive relations mix electric and magnetic fields

$$\mathbf{D}(\mathbf{r}, \omega) = \epsilon(\omega)\mathbf{E}(\mathbf{r}, \omega) - i\kappa(\omega)\mathbf{H}(\mathbf{r}, \omega)$$

$$\mathbf{B}(\mathbf{r}, \omega) = i\kappa(\omega)\mathbf{E}(\mathbf{r}, \omega) + \mu(\omega)\mathbf{H}(\mathbf{r}, \omega)$$

dispersive chirality:  $\kappa(\omega) = \frac{\omega_k \omega}{\omega^2 - \omega_{\kappa R}^2 + i\gamma_k \omega}$

# Repulsion and chiral MMs

In chiral MMs the reflection matrix is non-diagonal (mixing of E and H fields).


The integrand of the Casimir-Lifshitz force between two identical chiral MMs has the form:

$$F = \frac{(r_{ss}^2 + r_{pp}^2 - 2r_{sp}^2)e^{-2Kd} - 2(r_{sp}^2 + r_{ss}r_{pp})^2e^{-4Kd}}{1 - (r_{ss}^2 + r_{pp}^2 - 2r_{sp}^2)e^{-2Kd} + (r_{sp}^2 + r_{ss}r_{pp})^2e^{-4Kd}}$$

One might achieve repulsive Casimir forces with strong chirality (i.e., large values of  $r_{sp}$ )

Same-chirality (SC) materials: repulsion

Opposite-chirality (OC) materials: repulsion




# Beyond EMA


Everything discussed so far is based on the assumption that the **effective medium approximation** (EMA) holds. We recall that this amounts to treating the MM in the “long-wavelength approximation”, i.e., field wavelengths much larger than the typical size of the unit cell of the MM.

How to calculate Casimir forces when EMA does not hold?


Can one trust predictions of Casimir repulsion with MMs based on EMA?



# Exact method: Scattering theory



The Casimir force still may be described in terms of reflections (scattering theory)



Symbolically, we may write the Casimir energy as

$$\frac{E(d)}{A} = \hbar \int_0^\infty \frac{d\xi}{2\pi} \log \det [1 - \mathcal{R}_1 e^{-\kappa d} \mathcal{R}_2 e^{-\kappa d}]$$

where  $\mathcal{R}_i \equiv \mathcal{R}_i(\mathbf{k}_{\parallel}, \mathbf{k}'_{\parallel}, p, p', i\xi)$

# Solving for the reflection matrix

The reflection matrix can be obtained with standard methods of numerical electromagnetism. One way is to solve Maxwell equations for the transverse fields

$$\begin{aligned}-ik \frac{\partial \mathbf{E}_t}{\partial z} &= \nabla_t [\chi \hat{e}_3 \cdot \nabla \times \mathbf{H}_t] - k^2 \mu \hat{e}_3 \times \mathbf{H}_t \\ -ik \frac{\partial \mathbf{H}_t}{\partial z} &= -\nabla_t [\zeta \hat{e}_3 \cdot \nabla \times \mathbf{E}_t] + k^2 \epsilon \hat{e}_3 \times \mathbf{E}_t\end{aligned}$$

Assuming a two-dimensional periodic structure, we have

$$\mathbf{E}_t(x, y) = e^{i\mathbf{k} \cdot \mathbf{r}} \sum_{m,n} \mathcal{E}_{m,n} \exp \left[ i \frac{2\pi n}{L_x} x + i \frac{2\pi m}{L_y} y \right]$$

$$\mathbf{H}_t(x, y) = e^{i\mathbf{k} \cdot \mathbf{r}} \sum_{m,n} \mathcal{H}_{m,n} \exp \left[ i \frac{2\pi n}{L_x} x + i \frac{2\pi m}{L_y} y \right]$$

where

$$\epsilon(x, y) = \sum_{m,n} \epsilon_{m,n} \exp \left[ i \frac{2\pi n}{L_x} x + i \frac{2\pi m}{L_y} y \right]$$

$$\mu(x, y) = \sum_{m,n} \mu_{m,n} \exp \left[ i \frac{2\pi n}{L_x} x + i \frac{2\pi m}{L_y} y \right]$$



# Exact reflection matrix

One can then write the equations for the transverse fields as



$$\boxed{-ik \frac{\partial \Psi_{m'n'}}{\partial z} = \sum_{mn} H_{m'n',mn} \Psi_{mn}} \quad \Psi_{mn} = \begin{bmatrix} \mathcal{E}_{mn}^x \\ \mathcal{E}_{mn}^y \\ \mathcal{H}_{mn}^x \\ \mathcal{E}_{mn}^y \end{bmatrix} = \begin{bmatrix} \psi_{mn}^1 \\ \psi_{mn}^2 \\ \psi_{mn}^3 \\ \psi_{mn}^4 \end{bmatrix}$$

Here  $H$  is a complicated matrix, that encapsulated the coupling of modes in the periodic structure.

By numerically solving this equation and imposing the proper boundary conditions of the field on the vacuum-metamaterial interphase (RCWA or S-matrix techniques), one can find the reflection matrix of the MM.

# 2D periodic structures - finite T

Casimir force between a Au sphere and Si pillars/grating/membrane @ T=300 K




$R = 50\mu\text{m}$   
period = 400 nm  
depth = 1070 nm


Davids, Intravaia, Rosa, DD, arXiv:1008.3580

Related techniques in 1D gratings at T=0K developed by Guérout, Lambrecht, Chan, et al. arXiv:1009.3487

# Chiral metamaterials - $T=0$



Effect of inhomogeneity across displacements  $x$



Total force relative to parallel metal plates

“repulsive” effect (force reduction) of chirality is **one ten-thousandth** of this!

## Conclusion

In the regime where the chiral metamaterial limit is valid, the effect is **too small** to be observable.

# Remarks: MMs and Casimir

- Metamaterials offer an interesting possibility for Casimir force manipulation: engineered optical response, (maybe) broadband, dynamic control.
- Several proposals for MM-based Casimir force use effective medium approximation. Their predictions have to be carefully checked since EMA breaks down for electromagnetic fluctuations with wavelengths comparable to metamaterial feature sizes.
- Casimir repulsion in vacuum-separated metallic/dielectric metamaterial structures seems hard to achieve. It is certainly impossible in geometries that are effectively one-dimensional (Casimir stability considerations).

## Casimir-Lifshitz theory and magneto-dielectric metamaterials

Henkel and Joulain, EPL **72**, 929 (2005)

Leonhardt and Philbin, NJP **9**, 254 (2007)

da Rosa, DD, Milonni, PRL **100**, 183602 (2008)

da Rosa, DD, Milonni, PRA **78**, 032117 (2008)

## Numerical techniques for Casimir interactions with metamaterials


McCauley et al, PRB **82**, 165108 (2010)

Davids, Intravaia, Rosa, DD, arXiv:1008.3580



# New results (under review)



## First observation of the thermal Casimir force



Sushkov (Yale), Kim (Seattle), DD (LANL), and Lamoreaux (Yale)

# Torsional Pendulum Set-up

## \* Upgrade of Lamoreaux's 1997 experiment




An imbalance in capacitance is amplified and sent to a phase sensitive detector (PSD), which generates error signals.

A proportional integro-differential (PID) controller provides a feedback correction voltage  $S_{\text{PID}}(d, V_a)$  to the compensator plates, restoring equilibrium.

$$F \propto (S_{\text{PID}} + 9V)^2 \approx (9V)^2 + 2S_{\text{PID}} \times 9V$$

The correction voltage is the physical observable, and it is proportional to the force between the Casimir plates





# Typical Casimir Measurement

$$S_{\text{PID}}(d, V_a) = S_{\text{dc}}(d \rightarrow \infty) + S_a(d, V_a) + S_r(d)$$

force-free component of  
signal at large separations

electrostatic signal in  
response to an applied  
external voltage

residual signal due to  
distance-dependent  
forces, e.g. Casimir

The electrostatic signal between the spherical lens and the plate, in PFA ( $d \ll R$ ), is

$$S_a(d, V_a) = \pi \epsilon_0 R (V_a - V_m)^2 / \beta d \quad \beta \text{ force-voltage conversion factor}$$


This signal is minimized ( $S_a = 0$ ) when  $V_a = V_m$ , and the electrostatic minimizing potential  $V_m$  is then defined to be the **contact potential** between the plates.

# “Parabola” measurements

## Calibration routine (Iannuzzi et al, PNAS 04)

A range of plate voltages  $V_a$  is applied, and at a given nominal absolute distance the response is fitted to a parabola

$$S_{\text{PID}}(d, V_a) = S_0 + k(V_a - V_m)^2$$



## Fitting parameters


- $k = k(d) \longrightarrow$  voltage-force calibration factor + absolute distance
- $V_m = V_m(d) \longrightarrow$  distance-dependent minimizing potential
- $S_0 = S_0(d) \longrightarrow$  force residuals: electrostatic + Casimir + non-Newtonian gravity + ...

This procedure is repeated at decremental distances, from 7  $\mu\text{m}$  down to 0.7  $\mu\text{m}$ , completing a single experimental run.

In the experiment  $V_m = V_m(d)$  is almost constant (0.2 mV variation in the whole range)

# Force Residuals

Residuals from Coulomb force obtained from the value of the PID signal at the minima of each parabola,




In the experiment, these force residuals are too large to be explained just by the Casimir-Lifshitz force between the Au plates.

# Electrostatic Patch Effects

## Sphere-plane geometry:

To compute the patch effect in the sphere-plane configuration we use PFA for the curvature effect ( $d \ll R$ ) but leave  $kd$  arbitrary



$$\nabla^2 V(x, y, z) = 0$$

$$F_{sp}(d) = 2\pi R \langle U_{pp}(d) \rangle = \frac{\epsilon_0 R}{16} \int_0^\infty dk \frac{k^2 e^{-kd}}{\sinh(kd)} [C_{1,k} + C_{2,k}]$$

$$V(z=0) = V_1(x, y)$$

## Different models to describe surface potential fluctuations:

•  $C_{1,k} = C_{2,k} = V_0^2$  for  $k_{\min} < k < k_{\max}$

$$F_{sp} = \frac{4\pi\epsilon_0 V_{\text{rms}}^2 R}{k_{\max}^2 - k_{\min}^2} \int_{k_{\min}}^{k_{\max}} dk \frac{k^2 e^{-kd}}{\sinh(kd)}$$

•  $\mathcal{R}(r) = \begin{cases} V_0^2 & \text{for } r \leq \lambda, \\ 0 & \text{for } r > \lambda. \end{cases}$

$$F_{sp} = 2\pi\epsilon_0 R \int_0^\infty du \, u \frac{J_1(u)}{e^{2ud/\lambda} - 1}$$

In the limit of large patches ( $kd \ll 1$ ):

$$F_{sp}(d) = \pi\epsilon_0 R \frac{V_{\text{rms}}^2}{d}$$


# Understanding elec. residuals

We fit the data for the *residual force at the minimizing potential* with a force of equal to Casimir + patch effect

$$F_r(d) = F_C(d) + \pi\epsilon_0 R \frac{V_{\text{rms}}^2}{d}$$

Drude, T=300K


Drude, T=0K



Plasma, T=300K

Plasma, T=0K

# Thermal Casimir force



$$F_C^{(T)}(\text{Drude}) = \frac{\xi(3)}{8} \frac{Rk_B T}{d^2}$$

Quality of fits:

Drude,  $T=300\text{K}$

Plasma,  $T=300\text{K}$

Drude,  $T=0\text{K}$

Plasma,  $T=0\text{K}$

$\chi_{\text{red}}^2$      $V_{\text{rms}}(\text{mV})$

1.04	5.4
32	3.0
23	4.0
43	3.6

# Remarks:

- Experiment rules out the plasma model in the separation range 0.7um to 7um, and confirms the Drude model

- Thermal correction to the Casimir force demonstrated.

- Electrostatic residuals modeled as due to large electrostatic patches

$$F_r^{\text{patches}} \propto R \frac{V_{\text{rms}}^2}{d} \quad (\lambda_P \gg d)$$

- Would be interesting to do Kelvin probe microscopy of the used samples. Must detect potential variations on the 0.1 mV range.